

# BOUNDING SURFACE ACTIONS ON HYPERBOLIC SPACES

JOSH BARNARD

**ABSTRACT.** We give a diameter bound for fundamental domains for isometric actions of the fundamental group of a closed orientable hyperbolic surface on a  $\delta$ -hyperbolic space, where the bound depends on the hyperbolicity constant  $\delta$ , the genus of the surface, and the injectivity radius of the action, which we assume to be strictly positive.

## 1. INTRODUCTION

If  $S$  is a closed hyperbolic surface with genus  $g$  and injectivity radius bounded below by  $\epsilon > 0$ , then the diameter of  $S$  is bounded above by some constant  $D$ , depending only on  $g$  and  $\epsilon$ . One way to prove this is as follows (cf. [Bon86, Lemma 1.10]). The total area of  $S$  is bounded in terms of the genus  $g$  and the curvature  $\kappa = -1$ . Now join any pair of points by a distance-realising path of length  $D$ . The injectivity radius assumption implies that such a path has an embedded product neighborhood with radius  $\epsilon/3$ , the area of which (by the negative curvature assumption) is at least  $D\epsilon/3$ . It follows that  $D$  is constrained by the global area bound.

This result can be rephrased so as to provide a bound for the diameter of a fundamental domain for the corresponding action of  $\pi_1(S)$  on  $\mathbf{H}^2$ . In this context, we seek a generalization to actions of  $\pi_1(S)$  on  $\delta$ -hyperbolic spaces. In place of area bounds, which are no longer available, we exploit the fact that a large diameter requires the existence of short loops in the quotient.

Suppose  $\Gamma$  is a spine on  $S$ ; i.e.,  $\Gamma$  is a finite connected graph embedded in  $S$  with single disk complement. We let  $\tilde{\Gamma}$  denote the lift of  $\Gamma$  to the universal cover  $\tilde{S}$  of  $S$ . Suppose  $\gamma: \tilde{\Gamma} \rightarrow X$  is an embedding into a length-space  $X$  that is equivariant with respect to some isometric action of  $\pi_1(S)$  on  $X$ . We obtain a pseudometric  $d_\gamma$  on  $\Gamma$  by declaring

$$d_\gamma(p, q) = \inf\{d_X(P, Q) \mid P \in \gamma(\tilde{p}), Q \in \gamma(\tilde{q})\}.$$

This will be a true metric when the injectivity radius of the action is strictly positive; i.e., when

$$\text{inj}(X, \pi_1(S)) = \inf_{1 \neq h \in \pi_1(S)} \left( \lim_{n \rightarrow \infty} \frac{d_X(x, h^n x)}{n} \right) \geq \epsilon > 0.$$

From this metric we obtain a notion of diameter, given by

$$\text{diam}_\gamma(\Gamma) = \max\{d_\gamma(a, b) \mid a, b \in \Gamma\}.$$

Our main result is the following.

---

*Date:* November 20, 2009.

*2000 Mathematics Subject Classification.* 20F65.

**Theorem.** *There is a constant  $D = D(\delta, g, \epsilon)$  with the following property: Suppose  $S$  is a closed, orientable surface of genus  $g$ , and suppose  $\pi_1(S)$  acts isometrically on a  $\delta$ -hyperbolic space  $X$  with  $\text{inj}(X, \pi_1(S)) \geq \epsilon > 0$ . Then there is a spine  $\Gamma$  embedded on  $S$  and a  $\pi_1(S)$ -equivariant embedding  $\gamma: \tilde{\Gamma} \rightarrow X$  so that  $\text{diam}_\gamma(\Gamma) \leq D$ .*

**Remark 1.1.** We originally proved (a slightly less general version of) this theorem in the appendix to [Bar04]. A slightly different formulation of this theorem says that given a fixed generating set for  $\pi_1(S)$ , there is some point  $x \in X$  that is translated a bounded distance by each element in that generating set. See [Bow07a] for more details. In [Bow07b] Bowditch shows how to adapt the proof here to apply also to surfaces with boundary, as well as to nonorientable surfaces. In [DF09] Dahmani and Fujiwara prove a version of Theorem 1, where a general one-ended subgroup of the mapping class group acts on the curve complex of the surface.

**Acknowledgements.** Thanks are due to Brian Bowditch for many helpful comments, as well as for motivating me to write this up properly. I am also extremely grateful to Daryl Cooper, who, among other things, suggested the use of carrier graphs, as used in [Whi00].

## 2. PRELIMINARIES

We first fix some notation that will hold for the rest of the paper. We let  $S$  denote a closed orientable surface of genus  $g$ , and we assume we have a fixed isometric action of  $\pi_1(S)$  on a complete geodesic path-metric space  $(X, d_X)$  with  $\delta$ -thin triangles. The injectivity radius of this action is assumed to be bounded below by  $\epsilon > 0$ .

Let  $\tilde{S}$  denote the universal cover of  $S$ , and for any subset  $Z \subset S$ , we let  $\tilde{Z}$  denote the full lift of  $Z$  in  $\tilde{S}$ . A *carrier graph* for the action of  $\pi_1(S)$  on  $X$  is a pair  $(\Gamma, \gamma)$ , where  $\Gamma$  is an embedded graph  $\Gamma \subseteq S$  carrying all of  $\pi_1(S)$  and  $\gamma: \tilde{\Gamma} \rightarrow X$  is a  $\pi_1(S)$ -equivariant map sending edges of  $\Gamma$  to rectifiable paths in  $X$ . Given a carrier graph  $(\Gamma, \gamma)$ , we obtain a metric  $d_\gamma$  as described above, and we let  $\ell_\gamma(\Gamma)$  denote the sum of the lengths of the edges of  $\Gamma$  measured in this way. We abuse notation by using  $d_\gamma$ ,  $\ell_\gamma$ , and  $\text{diam}_\gamma$  also for arbitrary subsets of  $S$ .

We say that  $(\Gamma, \gamma)$  is *minimal* if  $\ell_\gamma(\Gamma) = \inf\{\ell_\sigma(\Sigma)\}$ , where the infimum is taken over all carrier graphs  $(\Sigma, \sigma)$  for the given action. Note that if  $(\Gamma, \gamma)$  is a minimal carrier graph, then we may assume that  $\Gamma$  is a spine on  $S$ . For convenience we will assume all vertices of  $\Gamma$  are at least trivalent.

**Remark 2.1.** A priori this infimum may not be achieved, in which case one needs to work with *almost minimal* carrier graphs, and then let the error term go to zero. For convenience of exposition, we will assume that a minimal carrier graph exists for the given action.

## 3. PROOF

We say a set  $Z \subseteq X$  has *controlled diameter* if it is contained in the  $\pi_1(S)$ -orbit of a bounded number of sets, each with bounded diameter, where all bounds depend on  $\delta$ ,  $g$ , and  $\epsilon$ . We verify the theorem by showing that  $\gamma(\tilde{\Gamma})$  has controlled diameter.

Fix a minimal spine  $(\Gamma, \gamma)$  and let  $\Delta$  denote the closure in  $\tilde{S}$  of a single lift of the complement  $S - \Gamma$ . A simple Euler characteristic argument shows that the number of edges of any spine is bounded above by  $6g - 3$ , so it follows from the minimality assumption that  $\gamma(\partial\Delta)$  is a geodesic  $n$ -gon in  $X$ , where  $n \leq 12g - 6$ . Note that the action of  $\pi_1(S)$  on  $X$  identifies sides of  $\gamma(\partial\Delta)$  in pairs so that the quotient is  $\Gamma$ .

Let  $\Gamma^{(0)}$  denote the vertices of  $\Gamma$ , and define  $B_0$  to be the  $n\delta$ -neighborhood of  $\gamma(\tilde{\Gamma}^{(0)})$  in  $X$ . Note that  $B_0$  has controlled diameter, as the number of vertices of  $\Gamma$  is bounded in terms of  $g$ . Suppose there is some point  $\tilde{s} \in \partial\Delta$  with  $\gamma(\tilde{s}) \notin B_0$ . Because  $\gamma(\partial\Delta)$  is a geodesic  $n$ -gon in a  $\delta$ -hyperbolic space, there is a point  $\tilde{s}'$  on  $\partial\Delta$  with  $d_X(\gamma(\tilde{s}), \gamma(\tilde{s}')) \leq n\delta$ . In fact, we may join  $\tilde{s}$  to  $\tilde{s}'$  by a path  $\tilde{\sigma}$  properly embedded in  $\Delta$  and extend  $\gamma$  to  $\tilde{\sigma}$  so that  $\ell_\gamma(\tilde{\sigma}) \leq n\delta$ .

**Claim.** *The two points  $\tilde{s}$  and  $\tilde{s}'$  project into  $S$  to the same edge  $e$  of  $\Gamma$ . Moreover, the  $\gamma$ -length of that portion of  $e$  bounded by the projections  $s, s' \in \Gamma$  has length no more than  $n\delta$ .*

*Proof.* Suppose first that  $s$  and  $s'$  lie on distinct edges  $e$  and  $e'$  of  $\Gamma$ , and let  $\sigma$  be the projection of  $\tilde{\sigma}$  into  $S$ . We will obtain a contradiction by constructing a carrier graph  $(\Gamma', \gamma')$  with  $\ell_{\gamma'}(\Gamma') < \ell_\gamma(\Gamma)$ . We construct  $\Gamma'$  from  $\Gamma$  by removing a component of  $e - s$  and inserting the path  $\sigma$ , and then  $\gamma'$  is simply the appropriate restriction of  $\gamma$  (recall that we had extended  $\gamma$  to  $\tilde{\sigma}$ ).

Because the complement of  $\Gamma$  is a disk, the complement of  $\Gamma \cup \sigma$  is two disks. It follows that  $\Gamma'$  has a single disk complement and thus carries  $\pi_1(S)$ . Because  $\gamma(\tilde{s})$  lies outside  $B_0$ , the segment of  $\Gamma$  deleted in forming  $\Gamma'$  has  $\gamma$ -length greater than  $n\delta$ , while  $\sigma$  has  $\gamma'$ -length less than  $n\delta$ . This contradicts the assumed minimality of  $(\Gamma, \gamma)$ . We deduce that  $e = e'$ .




FIGURE 1. Two ways to shorten a carrier graph (the lengths suggested by the pictures are  $\gamma$ -lengths).

Now suppose the  $\gamma$ -length of the portion of  $e$  between  $s$  and  $s'$  is greater than  $n\delta$ . We will again arrive at a contradiction by constructing a shorter carrier graph. This time we construct  $\Gamma'$  by deleting that portion of  $e$  between  $s$  and  $s'$  and inserting  $\sigma$ , and define  $\gamma'$  as before. The result follows.  $\square$

It follows from this claim that  $\sigma$  and that portion of  $e$  between  $s$  and  $s'$  together form a simple loop  $\beta_1$  on  $S$ . This loop is homotopically essential, as it does not lift to a loop in  $\tilde{S}$ .


Also note that the claim implies that  $\ell_\gamma(\beta_1) \leq 2n\delta$ . Thus from the existence of a point on  $\partial\Delta$  mapping outside  $B_0$ , we construct an essential closed curve on  $S$  with  $\gamma$ -length no more than  $2n\delta$ .

The plan now is to enlarge  $B_0$  to  $B_1$ , which will include a neighborhood of the  $\pi_1(S)$ -orbit of  $\gamma(\tilde{\beta}_1)$ . We then proceed as before, showing that the existence of a point of  $\gamma(\partial\Delta)$  outside  $B_1$  provides another simple closed curve on  $S$  with short  $\gamma$ -length. Moreover, this curve will be disjoint from and not freely-homotopic to the previous one. For all of this we need the following lemma.

**Lemma 3.1.** *There is a constant  $K$ , depending only on  $\delta$ ,  $n$ , and  $\epsilon$ , so that if  $x$  and  $x'$  are two points in  $X$  with the property that  $d_X(x, hx) \leq 2n\delta$  and  $d_X(x', hx') \leq 2n\delta$  for some  $h \in \pi_1(S)$ , then the Hausdorff distance between  $\{h^n x\}$  and  $\{h^n x'\}$  is bounded above by  $K$ .*

*Proof.* Here is a sketch of the proof. Because of the injectivity radius lower bound, every element of  $\pi_1(S)$  has a quasi-axis in  $X$ ; i.e., for each  $h \in \pi_1(S)$  there is a quasigeodesic  $A_h$  moved by the element a bounded Hausdorff distance from itself, where the bounds all depend only on  $\delta$ . We claim that the farther a point is from this quasixaxis, the greater the distance it is moved by  $h$ . Thus an upper bound on the distance a point is moved by  $h$  gives an upper bound of the distance from the point to the quasixaxis, where the bound depends on  $\delta$  as well as the stable translation length of  $h$ , which in turn is bounded below by  $\epsilon$ . In particular, two such points are boundedly Hausdorff close to the quasixaxis, and hence to one another.

To prove the claim, pick  $x \in X$ , and suppose  $d_X(x, A_h) = d_X(x, a) = L$  for some point  $a$  on the quasixaxis  $A_h$ . Fix  $m = \lceil 2\delta/\epsilon \rceil$ , so that  $d_X(a, h^m x) \geq 2\delta$ . Let  $\alpha$  denote a geodesic segment joining  $x$  to  $h^m x$ . For sufficiently large  $L$ , we may choose  $p$  on a geodesic segment joining  $x$  to  $a$  so that  $\delta < d_X(p, a) < 2\delta$ . Now suppose  $A_h$  is actually geodesic. Then we also have  $\delta < d_X(p, A_h) < 2\delta$ , so by the thin triangles condition applied to the quadrilateral  $(a, x, h^m x, h^m a)$ , there is some point  $p' \in \alpha$  so that  $d_X(p, p') \leq \delta$ . By the triangle inequality,



we have  $d_X(x, p') \geq L - 3\delta$ . We similarly find  $q'$  with  $d_X(h^m x, q') \geq L - 3\delta$ . By choice of  $m$ , the portion of  $\alpha$  joining  $x$  to  $p'$  does not intersect the portion joining  $h^m x$  to  $q'$ . It follows that  $d_X(x, h^m x) \geq 2L - 6\delta$ . Dividing by  $2\delta/\epsilon$ , we find that  $d_X(x, hx) \geq \frac{L\epsilon}{\delta} - 3\epsilon$ , so that

$h$ -translation distance is an unboundedly increasing function of  $L$ , as claimed. When  $A_h$  is not geodesic, we apply the argument to a geodesic segment  $\overline{A}$  joining  $a$  to  $h^m a$  and use the fact that the corresponding portion of  $A_h$  lies within a bounded Hausdorff neighborhood of  $\overline{A}$ .  $\square$

Using this constant  $K$ , we define  $B_1$  to be the union of  $B_0$  with the  $\max\{K, 4n\delta\}$ -neighborhood of  $\gamma(\tilde{\beta}_1)$ . Note that  $B_1$  has controlled diameter. This is because the  $\gamma$ -length of  $\beta_1$  is bounded appropriately, as is the diameter of the neighborhood taken.

For general  $j \geq 1$ , suppose  $B_{j-1}$  has been defined, and suppose  $\gamma(\partial\Delta) \not\subset B_{j-1}$ . Then we may use a point on  $\partial\Delta$  mapping outside  $B_{j-1}$  to construct a simple closed curve  $\beta_j$  on  $S$  with  $\gamma$ -length (for  $\gamma$  extended as before) no more than  $2n\delta$ . We then inductively define  $B_j$  to be the union of  $B_{j-1}$  with the  $\max\{K, 4n\delta\}$ -neighborhood of  $\gamma(\tilde{\beta}_j)$ . Note that  $B_j$  has controlled diameter.

We claim now that  $\beta_j \cap \beta_i = \emptyset$  for  $i \neq j$ , and that  $\beta_j$  and  $\beta_i$  are not freely homotopic. For the first part, if the two curves intersected in some point  $p$ , then because each of  $\beta_j$  and  $\beta_i$  has  $\gamma$ -length no more than  $2n\delta$ , one could construct a path from any point on  $\beta_j$  to any point on  $\beta_i$ , through  $p$ , with total  $\gamma$ -length no more than  $4n\delta$ . This contradicts the fact at least one point on  $\gamma(\tilde{\beta}_j)$  lies outside the  $4n\delta$ -neighborhood of  $\gamma(\tilde{\beta}_i)$  for  $i < j$ .

For the second part, because at least one point of  $\gamma(\tilde{\beta}_j)$  lies outside the  $K$ -neighborhood of all previous  $\gamma(\tilde{\beta}_i)$ , the lemma above implies that  $\beta_j$  cannot be freely homotopic to any  $\beta_i$  for  $i < j$ .

The result follows when we note that the maximum number of non-parallel disjoint simple closed curves  $\beta_j$  one may find on  $S$  is  $2g - 1$ , so that  $\gamma(\partial\Delta)$  must be contained in  $B_k$  for some  $k \leq 2g - 1$ . But the  $\pi_1(S)$ -orbit of  $\gamma(\partial\Delta)$  is all of  $\gamma(\tilde{\Gamma})$ . Thus  $\gamma(\tilde{\Gamma})$  is contained in the set  $B_k$ , which has controlled diameter. The result follows.

## REFERENCES

- [Bar04] J. Barnard. *Ends of word-hyperbolic three-manifolds*. PhD thesis, Univ. of California, Santa Barbara, 2004.
- [Bon86] F. Bonahon. Bouts des variétés hyperboliques de dimension 3. *Ann. Math.*, 124:71–158, 1986.
- [Bow07a] B. H. Bowditch. Atoroidal surface bundles over surfaces. *Preprint*, 2007. Available at <http://www.warwick.ac.uk/~masgak/>.
- [Bow07b] B. H. Bowditch. One-ended subgroups of mapping class groups. *Preprint*, 2007. Available at <http://www.warwick.ac.uk/~masgak/>.
- [DF09] François Dahmani and Koji Fujiwara. Copies of one-ended groups in mapping class groups. *Groups Geom. Dyn.*, 3(3):359–377, 2009.
- [Whi00] M. White. *Some bounds for closed hyperbolic 3-manifolds*. PhD thesis, Univ. of California, Santa Barbara, 2000.

DEPT. OF MATHEMATICS & STATISTICS, UNIVERSITY OF SOUTH ALABAMA, MOBILE, AL 36688  
 E-mail address: [jbarnard@jaguar1.usouthal.edu](mailto:jbarnard@jaguar1.usouthal.edu)